

Aczél–Chebyshev Type Inequality for Positive Linear Functions¹

Xie-Hua Sun²

metadata, citation and similar papers at core.ac.uk

Submitted by B. G. Pachpatte

Received October 27, 1998

An Aczél–Chebyshev type inequality and some exact inequalities like the ones of Karamata for positive linear functionals are established. A counterexample on Chebyshev type inequality is given. © 2000 Academic Press

1. INTRODUCTION

Denote $\mathcal{R} = (-\infty, +\infty)$ and $\mathcal{R}^+ = [0, +\infty)$.

Let \mathcal{E} be a nonempty set on \mathcal{R} and let \mathcal{L} be a class of nonnegative functions $f: \mathcal{E} \rightarrow \mathcal{R}^+$ and $\mathcal{F} \supset \mathcal{L}$ the class of functions $f: \mathcal{E} \rightarrow \mathcal{R}$ on which the functionals A are defined. We shall consider functionals $A: \mathcal{F} \rightarrow \mathcal{R}$ which satisfy the following conditions.

- (1) $f \in \mathcal{L} \Rightarrow A(f) \geq 0$.
- (2) $f \in \mathcal{F}, \lambda \in \mathcal{R} \Rightarrow \lambda f \in \mathcal{F}, A(\lambda f) = \lambda A(f)$.
- (3) $1 \in \mathcal{F}, A(1) = 1$.
- (4) $f, g \in \mathcal{F} \Rightarrow f + g \in \mathcal{F}, A(f + g) = A(f) + A(g)$.

If a functional A satisfies conditions (1)–(4), we say A is a *positive linear functional*.

Pečarić [5] obtained generalizations of fundamental mean inequality, Hölder's and Minkowski's, and their converse inequality. In the same paper, Pečarić also considered the functional $p(f) = (f_0^r - A(f^r))^{1/r}$ and gave some results which generalized the well-known Aczél, Popoviciu, and Bellman inequalities.

¹ This work was supported by Zhejiang Provincial Natural Science Foundation of China.

² E-mail address: xhsun@mail.hz.zj.cn.

In this paper, we shall consider the functional $p(f) = f_0 - A(f)$. In Section 2, we will establish an Aczél–Chebyshev type inequality for monotone functions $f(x)$ and $g(x)$ in the opposite sense. It is well known that if $f(x)$ and $g(x)$ are monotone in the same sense, we will have the reverse Chebyshev inequality (see [4]). Unfortunately, in this case, we cannot obtain the reverse Aczél–Chebyshev type inequality for $f(x)$ and $g(x)$ monotone in the same sense. In Section 3, we will give a counterexample which shows this so-called reverse inequality cannot hold. In Section 4, we will establish some exact inequalities like the ones of Karamata (see [2, 3]).

2. ACZÉL–CHEBYSHEV TYPE INEQUALITY

In order to prove our main result, we need the following lemma, which is a generalization of Chebyshev inequality for a positive linear functional.

LEMMA 2.1 (Chebyshev type inequality). *Let A be a positive linear functional with $A(1) = 1$. If $f(x)$ and $g(x)$ are monotone in the same sense, then*

$$A(fg) \geq A(f)A(g). \quad (2.1)$$

Moreover, if $f(x)$ and $g(x)$ are monotone in the opposite sense, then

$$A(fg) \leq A(f)A(g). \quad (2.2)$$

Proof. Assume f and g are increasing on \mathcal{E} . Then we have

$$\inf_{x \in \mathcal{E}} g(x) \leq A(g) \leq \sup_{x \in \mathcal{E}} g(x).$$

Therefore, there exists a $\xi \in \mathcal{E}$ satisfying

$$g(x) \geq A(g) \quad \text{if } x \in \mathcal{E}, \quad x \geq \xi,$$

and

$$g(x) \leq A(g) \quad \text{if } x \in \mathcal{E}, \quad x < \xi,$$

or satisfying

$$g(x) \geq A(g) \quad \text{if } x \in \mathcal{E}, \quad x > \xi,$$

and

$$g(x) \leq A(g) \quad \text{if } x \in \mathcal{E}, \quad x \leq \xi.$$

Hence, we have, for all $x \in \mathcal{E}$,

$$f(x)(g(x) - A(g)) \geq f(\xi)(g(x) - A(g)). \quad (2.3)$$

Acting by A on the two sides of Eqs. (2.3) yields

$$\begin{aligned} A(fg) - A(f)A(g) &= A(f(g - A(g))) \\ &\geq f(\xi)A(g - A(g)) = 0. \end{aligned}$$

For the other cases, the proof of Eq. (2.1) is similar. We omit the detail.

THEOREM 2.1 (Aczél–Chebyshev type inequality). *Let A be a positive linear functional with $A(1) = 1$, and let f_0, g_0 be two nonnegative constants. If $f(x)$ and $g(x)$ are monotone in the opposite sense and $A(f) \geq 0, A(g) \geq 0, f_0 - A(f) \geq 0, g_0 - A(g) \geq 0$, then*

$$f_0 g_0 - A(fg) \geq (f_0 - A(f))(g_0 - A(g)). \quad (2.4)$$

Proof. If $f_0 = 0$ or $g_0 = 0$, then Eq. (2.4) follows from Eq. (2.2). Now assume $f_0 > 0$, and $g_0 > 0$. It is clear that Eq. (2.4) is equivalent to

$$1 - A(fg) \geq (1 - A(f))(1 - A(g)), \quad (2.5)$$

if we substitute f by f/f_0 , g by g/g_0 in Eq. (2.4), respectively. And Eq. (2.5) is equivalent to the following inequality:

$$A(f)(1 - A(g)) + A(g) - A(fg) \geq 0. \quad (2.6)$$

Finally, Eq. (2.6) is true since

$$A(f)(1 - A(g)) \geq 0,$$

and, applying Eq. (2.2),

$$A(g) - A(fg) \geq A(g) - A(f)A(g) = A(g)(1 - A(f)) \geq 0.$$

Remark. The conditions $A(f) \geq 0$ and $A(g) \geq 0$ in Theorem 2.1 are necessary. In fact, if we set

$$A(f) = \int_0^1 f(x) dx \quad (2.7)$$

and

$$f(x) = \begin{cases} -1, & x \in [0, \frac{1}{2}), \\ 0, & x \in [\frac{1}{2}, 1], \end{cases} \quad g(x) = \begin{cases} \frac{1}{10}, & x \in [0, \frac{1}{2}), \\ \frac{1}{20}, & x \in [\frac{1}{2}, 1], \end{cases}$$

then $f(x)$ and $g(x)$ are monotone in the opposite sense and satisfy $A(f) < 0$ and $A(g) \geq 0$. But we have

$$1 - A(fg) = 1 + \frac{1}{20} < (1 + \frac{1}{2})(1 - \frac{3}{40}) = (1 - A(f))(1 - A(g)).$$

3. A COUNTEREXAMPLE ON ACZÉL-CHEBYSHEV TYPE INEQUALITY

Now we give a counterexample which shows that if $f(x)$ and $g(x)$ are monotone in the same sense, generally, we cannot obtain so-called Aczél–Chebyshev type inequality.

Define the positive linear functional A as in Eq. (2.7). Obviously, the functional A satisfies condition $A(1) = 1$.

Let $0 \leq m \leq 10$ and

$$f(x) = g(x) = \begin{cases} m, & x \in [0, \frac{1}{10}], \\ 0, & x \in (\frac{1}{10}, 1]. \end{cases}$$

We have

$$\begin{aligned} (1 - A(f))(1 - A(g)) &= \left(1 - \int_0^1 f(x) dx\right) \left(1 - \int_0^1 g(x) dx\right) \\ &= \left(1 - \int_0^{1/10} m dx\right)^2 \\ &= \left(1 - \frac{m}{10}\right)^2 \end{aligned}$$

and

$$1 - A(fg) = 1 - \int_0^1 f(x)g(x) dx = 1 - \frac{m^2}{10}.$$

If $m = 3$, we have

$$\left(1 - \int_0^1 f(x) dx\right) \left(1 - \int_0^1 g(x) dx\right) > 1 - \int_0^1 f(x)g(x) dx.$$

If $m = \frac{1}{2}$, we obtain

$$\left(1 - \int_0^1 f(x) dx\right) \left(1 - \int_0^1 g(x) dx\right) < 1 - \int_0^1 f(x)g(x) dx.$$

This example shows that we cannot obtain the inequality

$$f_0 g_0 - A(fg) \leq (f_0 - A(f))(g_0 - A(g))$$

if $f(x)$ and $g(x)$ are monotone in the same sense. But we have the following

PROPOSITION 3.1. *Let A be a positive linear functional with $A(1) = 1$. If $0 \leq f(x) \leq f_0$ or $0 \leq g(x) \leq g_0$, we have*

$$(f_0 - A(f))(g_0 - A(g)) \leq f_0 g_0 - A(fg). \quad (3.1)$$

Proof. If $f_0 = 0$ or $g_0 = 0$, the inequality (3.1) is trivial. Now assume $f_0 > 0$, $g_0 > 0$. Then (3.1) is equivalent to the inequality

$$(1 - A(f))(1 - A(g)) \leq 1 - A(fg),$$

with $0 \leq f(x) \leq 1$ and $0 \leq g(x) \leq 1$. Then we have

$$1 - A(fg) \geq 1 - A(g) \geq (1 - A(f))(1 - A(g)).$$

■

4. SOME EXACT INEQUALITIES

We say a functional A is *quasiconvex* (see [1]) if for $f, g \in \mathcal{F}$ and $0 < \alpha < 1$, then

$$A(\alpha f + (1 - \alpha)g) \leq \max\{A(f), A(g)\}.$$

We say A is *quasiconcave* if

$$A(\alpha f + (1 - \alpha)g) \geq \min\{A(f), A(g)\}.$$

A functional $A(f)$ is said to be *continuous* for $f \in \mathcal{F}$ if, for arbitrary small $\epsilon > 0$, there exists a $\delta > 0$ such that

$$|A(g) - A(f)| < \epsilon,$$

provided $g \in \mathcal{F}$ and

$$\|f - g\|_C < \delta.$$

If $A(f)$ is continuous for every $f \in \mathcal{S} \subset \mathcal{F}$, we say $A(f)$ is *continuous* on the set \mathcal{S} . Obviously, every positive linear functional is continuous.

Denote the characteristic function on the set Q by

$$X_Q(x) = \begin{cases} 1, & x \in Q, \\ 0, & x \notin Q. \end{cases}$$

We say $f(x)$ is a *step function* on \mathcal{E} if

$$f(x) = \sum_{i=1}^n \lambda_i X_{Q_i}, \quad \text{for some } n \in \mathcal{N},$$

where λ_i ($i = 1, \dots, n$) are constants and

$$Q_i \subset \mathcal{E}, \quad \bigcup_{i=1}^n Q_i = \mathcal{E} \quad \text{and} \quad Q_i \cap Q_j = \emptyset \quad (i \neq j).$$

Let A be a positive linear functional with $A(1) = 1$. Define the functional

$$F(f, g) = \frac{f_0 g_0 - A(fg)}{(f_0 - A(f))(g_0 - A(g))},$$

for f and g satisfying $f_0 - A(f) > 0$ and $g_0 - A(g) > 0$.

Define the following classes of functions

$$B(m, M) = \{f(x) \mid 0 < m \leq f(x) \leq M, \forall x \in \mathcal{E}\},$$

$$B^*(m, M) = \left\{ \phi(x) \mid \phi(x) = \sum_{i=1}^n \lambda_i X_{Q_i}(x) \text{ is a step function,} \right. \\ \left. \lambda_i = m \text{ or } M \right\},$$

$$BI(m_1, M_1) = \{f(x) \mid m_1 \leq f(x) \leq M_1; f(x) \text{ is increasing on } \mathcal{E}\},$$

$$BD(m_2, M_2) = \{f(x) \mid m_2 \leq f(x) \leq M_2; f(x) \text{ is decreasing on } \mathcal{E}\}.$$

The purpose of this section is to calculate

$$\sup_{\substack{f \in B(m_1, M_1) \\ g \in B(m_2, M_2)}} F(f, g), \quad \inf_{\substack{f \in B(m_1, M_1) \\ g \in B(m_2, M_2)}} F(f, g),$$

and

$$\sup_{\substack{f \in BI(m_1, M_1) \\ g \in BD(m_2, M_2)}} F(f, g), \quad \inf_{\substack{f \in BI(m_1, M_1) \\ g \in BD(m_2, M_2)}} F(f, g).$$

In order to do that, we need the following lemmas.

LEMMA 4.1. *The functional $F(f, g)$ is quasiconvex and quasiconcave for f and g satisfying $f_0 - A(f) > 0$, $g_0 - A(g) > 0$.*

Proof. Assume $\alpha + \beta = 1$, $\alpha > 0$, $\beta > 0$. Since

$$F(\alpha f_1 + \beta f_2, g) = \frac{\alpha(f_0 - A(f_1))F(f_1, g) + \beta(f_0 - A(f_2))F(f_2, g)}{\alpha(f_0 - A(f_1)) + \beta(f_0 - A(f_2))},$$

then we have

$$\min\{F(f_1, g), F(f_2, g)\} \leq F(\alpha f_1 + \beta f_2, g) \leq \max\{F(f_1, g), F(f_2, g)\}.$$

Similarly, we can prove $F(f, g)$ is quasiconvex and quasiconcave for g . ■

LEMMA 4.2. Let A be a positive linear functional with $A(1) = 1$. If ϕ_1, ϕ_2 are step functions on \mathcal{E} ,

$$\begin{aligned}\phi_1(x) &= \sum_{i=1, i \neq k}^n \lambda_i X_{Q_i}(x) + mX_{Q_k}(x), \\ \phi_2(x) &= \sum_{i=1, i \neq k}^n \lambda_i X_{Q_i}(x) + MX_{Q_k}(x),\end{aligned}$$

where $\lambda_i = m$ or M and $0 \leq m < M \leq 1$, then for $0 \leq g(x) \leq 1$,

$$\frac{1 - A(\phi_1 g)}{1 - A(\phi_1)} < \frac{1 - A(\phi_2 g)}{1 - A(\phi_2)}. \quad (4.1)$$

Proof. Obviously, (4.1) is the inequality

$$\begin{aligned}& \frac{1 - A(g \sum_{i=1, i \neq k}^n \lambda_i X_{Q_i}) - mA(gX_{Q_k})}{1 - A(\sum_{i=1, i \neq k}^n \lambda_i X_{Q_i}) - mA(X_{Q_k})} \\ & < \frac{1 - A(g \sum_{i=1, i \neq k}^n \lambda_i X_{Q_i}) - MA(gX_{Q_k})}{1 - A(\sum_{i=1, i \neq k}^n \lambda_i X_{Q_i}) - MA(X_{Q_k})}.\end{aligned}$$

The above inequality is equivalent to

$$\begin{aligned}& A(gX_{Q_k}) \left(1 - A \left(\sum_{i \neq k} \lambda_i X_{Q_i} \right) \right) \\ & < A(X_{Q_k}) \left(1 - A \left(g \sum_{i \neq k} \lambda_i X_{Q_i} \right) \right).\end{aligned}$$

It is clear that the above inequality is true. ■

LEMMA 4.3. Suppose $f_0 > M_1$ and $g_0 > M_2$. The functional $F(f, g)$ is continuous on the set $f \in B(m_1, M_1)$ and $g \in B(m_2, M_2)$, respectively.

Proof. Since f and g are symmetric in the functional $F(f, g)$, it is sufficient to prove $F(f, g)$ is continuous on $f \in B(m_1, M_1)$. The continuity can be deduced easily from the equality

$$\begin{aligned} & F(f, g) - F(f^*, g) \\ &= \frac{1}{(f_0 - A(f))(f_0 - A(f^*))(g_0 - A(g))} \\ & \quad \times \{f_0 A((f^* - f)g) + f_0 g_0 A(f - f^*) \\ & \quad + A(f^* - f)A(f^*g) + A(f^*)A((f - f^*)g)\}. \end{aligned}$$

■

First we establish a general theorem.

THEOREM 4.1. *If functional $F(f)$ is continuous and quasiconvex on \mathcal{F} , then*

$$\sup_{f \in B(m, M)} F(f) = \sup_{\phi^* \in B^*(m, M)} F(\phi^*). \quad (4.2)$$

If functional $F(f)$ is continuous and quasiconcave on \mathcal{F} , then

$$\inf_{f \in B(m, M)} F(f) = \inf_{\phi^* \in B^*(m, M)} F(\phi^*). \quad (4.3)$$

Proof. Define step function on \mathcal{E}

$$\phi(x) = \sum_{i=1}^n \lambda_i X_{Q_i}(x),$$

where $m \leq \lambda_i \leq M$. If $m < \lambda_k < M$, then there exists an $\alpha \in (0, 1)$ such that

$$\lambda_k = \alpha m + (1 - \alpha)M.$$

Therefore,

$$\phi(x) = \alpha \phi_1(x) + (1 - \alpha) \phi_2(x),$$

where

$$\begin{aligned} \phi_1(x) &= \sum_{i=1, i \neq k}^n \lambda_i X_{Q_i}(x) + m X_{Q_k}(x), \\ \phi_2(x) &= \sum_{i=1, i \neq k}^n \lambda_i X_{Q_i}(x) + M X_{Q_k}(x). \end{aligned}$$

Since $F(f)$ is quasiconvex, then

$$F(\phi) \leq \max\{F(\phi_1), F(\phi_2)\}.$$

Using the similar method for $i = 1, \dots, n$, we can obtain

$$F(\phi) \leq F(\phi^*),$$

where ϕ^* is a step function satisfying $\lambda_i = m$ or M ($i = 1, \dots, n$). It is clear that for any measurable function $f(x)$ with $m \leq f(x) \leq M$ in Lebesgue sense and for arbitrary $\delta > 0$, there exists a step function $\phi(x)$ satisfying

$$\|f - \phi\|_C \leq \delta.$$

Then from the continuity of $F(f)$, we deduce

$$|F(f) - F(\phi)| < \epsilon.$$

Hence,

$$F(f) \leq F(\phi) + \epsilon \leq \epsilon + \sup_{\phi^* \in B^*(m, M)} F(\phi^*).$$

Since ϵ is arbitrarily small,

$$\sup_{f \in B(m, M)} F(f) \leq \sup_{\phi^* \in B^*(m, M)} F(\phi^*).$$

The inequality

$$\sup_{\phi^* \in B^*(m, M)} F(\phi^*) \leq \sup_{f \in B(m, M)} F(f)$$

is obvious. Therefore, Eq. (4.2) holds.

The proof of Eq. (4.3) is similar. ■

THEOREM 4.2. Suppose $0 \leq m_1 < M_1 < f_0$ and $0 \leq m_2 < M_2 < g_0$. The equalities

$$\sup_{\substack{f \in BI(m_1, M_1) \\ g \in BD(m_2, M_2)}} F(f, g) = \sup_{\substack{f \in B(m_1, M_1) \\ g \in B(m_2, M_2)}} F(f, g) = \frac{f_0 g_0 - M_1 M_2}{(f_0 - M_1)(g_0 - M_2)} \quad (4.4)$$

and

$$\inf_{\substack{f \in BI(m_1, M_1) \\ g \in BD(m_2, M_2)}} F(f, g) = \inf_{\substack{f \in B(m_1, M_1) \\ g \in B(m_2, M_2)}} F(f, g) = \frac{f_0 g_0 - m_1 m_2}{(f_0 - m_1)(g_0 - m_2)} \quad (4.5)$$

hold.

Proof. Applying Lemmas 4.1 and 4.2 and Theorem 4.1 yields

$$\sup_{\substack{f \in BI(m_1, M_1) \\ g \in BD(m_2, M_2)}} F(f, g) \leq \sup_{\substack{f \in B(m_1, M_1) \\ g \in B(m_2, M_2)}} F(f, g) = \sup_{\substack{f \in B^*(m_1, M_1) \\ g \in B^*(m_2, M_2)}} F(f, g). \quad (4.6)$$

Using Lemma 4.2, we get

$$\sup_{\substack{f \in B^*(m_1, M_1) \\ g \in B^*(m_2, M_2)}} F(f, g) \leq F(\bar{f}, \bar{g}) = \frac{f_0 g_0 - M_1 M_2}{(f_0 - M_1)(g_0 - M_2)}, \quad (4.7)$$

where $\bar{f}(x) \equiv M_1$, $\bar{g}(x) \equiv M_2$ ($\forall x \in \mathcal{E}$). Since $\bar{f} \in B^*(m_1, M_1)$ and $\bar{g} \in B^*(m_2, M_2)$, Eq. (4.4) follows from Eqs. (4.6) and (4.7).

As to Eq. (4.5), the proof is similar. ■

COROLLARY 4.2.1. *Let A be a positive linear functional with $A(1) = 1$. Suppose $f \in B(m_1, M_1)$ and $g \in B(m_2, M_2)$, where $0 \leq m_1 < M_1 < f_0$ and $0 \leq m_2 < M_2 < g_0$. Then the inequality*

$$\begin{aligned} K_*(f_0 - A(f))(g_0 - A(g)) \\ \leq f_0 g_0 - A(fg) \leq K^*(f_0 - A(f))(g_0 - A(g)), \end{aligned}$$

holds, where

$$K_* = \frac{f_0 g_0 - m_1 m_2}{(f_0 - m_1)(g_0 - m_2)}, \quad K^* = \frac{f_0 g_0 - M_1 M_2}{(f_0 - M_1)(g_0 - M_2)}.$$

Denote

$$F(f_1, f_2, \dots, f_n) = \frac{\prod_{k=1}^n (G_k - A(\prod_{k=1}^n f_k))}{\prod_{k=1}^n (G_k - A(f_k))},$$

where G_k ($k = 1, \dots, n$) are constants and $G_k - A(\prod_{k=1}^n f_k) > 0$ ($k = 1, \dots, n$).

COROLLARY 4.2.2. Suppose $f_k \in B(m_k, M_k)$ and $0 \leq m_k < M_k < G_k$ ($k = 1, \dots, n$). Then we have

$$\sup_{\substack{f_k \in B(m_k, M_k) \\ k=1, \dots, n}} F(f_1, \dots, f_n) = \frac{\prod_{k=1}^n G_k - \prod_{k=1}^n M_k}{\prod_{k=1}^n (G_k - M_k)},$$

and

$$\inf_{\substack{f_k \in B(m_k, M_k) \\ k=1, \dots, n}} F(f_1, \dots, f_n) = \frac{\prod_{k=1}^n G_k - \prod_{k=1}^n m_k}{\prod_{k=1}^n (G_k - m_k)}.$$

REFERENCES

1. M. Avriel, "Nonlinear Programming Analysis and Methods," Prentice-Hall, Englewood Cliffs, NJ, 1976.
2. J. Karamata, Inégalités relatives aux quotients et à la difference de $[fg]$ et $[f]g$, *Acad. Serbe Sci. Publ. Inst. Math.* **2** (1948), 131–145.
3. A. Lupaş, On two inequalities of Karamata, *Univ. Beograd. Publ. Elektrotehn. Fak. Ser. Mat. Fiz. No.* **602–633** (1978), 119–123.
4. D. S. Mitrinović, "Analytic Inequalities," Springer-Verlag, Berlin/Heidelberg/New York, 1970.
5. J. E. Pečarić, Generalization of the power means and their inequalities, *J. Math. Anal. Appl.* **161** (1991), 395–404.